

A COMPUTATION OF TIGHT CLOSURE IN DIAGONAL HYPERSURFACES

ANURAG K. SINGH

1. INTRODUCTION

The aim of this paper is to settle a question about the tight closure of the ideal (x^2, y^2, z^2) in the ring $R = K[X, Y, Z]/(X^3 + Y^3 + Z^3)$ where K is a field of prime characteristic $p \neq 3$. (Lower case letters denote the images of the corresponding variables.) M. McDermott has studied the tight closure of various irreducible ideals in R , and has established that $xyz \in (x^2, y^2, z^2)^*$ when $p < 200$, see [Mc]. The general case however existed as a classic example of the difficulty involved in tight closure computations, see also [Hu, Example 1.2]. We show that $xyz \in (x^2, y^2, z^2)^*$ in arbitrary prime characteristic p , and furthermore establish that $xyz \in (x^2, y^2, z^2)^F$ whenever R is not F-pure, i.e., when $p \equiv 2 \pmod{3}$. We move on to generalize these results to the diagonal hypersurfaces $R = K[X_1, \dots, X_n]/(X_1^n + \dots + X_n^n)$.

These issues relate to the question whether the tight closure I^* of an ideal I agrees with its plus closure, $I^+ = IR^+ \cap R$, where R is a domain over a field of characteristic p and R^+ is the integral closure of R in an algebraic closure of its fraction field. In this setting, we may think of the Frobenius closure of I as $I^F = IR^\infty \cap R$ where R^∞ is the extension of R obtained by adjoining p^e th roots of all nonzero elements of R for $e \in \mathbb{N}$. It is not difficult to see that $I^+ \subseteq I^*$, and equality in general is a formidable open question. It should be mentioned that in the case when I is an ideal generated by part of a system of parameters, the equality is a result of K. Smith, see [Sm]. In the above ring $R = K[X, Y, Z]/(X^3 + Y^3 + Z^3)$ where K is a field of characteristic $p \equiv 2 \pmod{3}$, if one could show that $I^* = I^F$ for an ideal I , a consequence of this would be $I^F \subseteq I^+ \subseteq I^* = I^F$, by which $I^+ = I^*$. McDermott does show that $I^* = I^F$ for large families of irreducible ideals and our result $xyz \in (x^2, y^2, z^2)^F$, we believe, fills in an interesting remaining case.

2. DEFINITIONS

Our main reference for the theory of tight closure is [HH]. We next recall some basic definitions.

Let R be a Noetherian ring of characteristic $p > 0$. We shall always use the letter e to denote a variable nonnegative integer, and q to denote the e th power of p , i.e., $q = p^e$. We shall denote by F , the Frobenius endomorphism of R , and by F^e , its e th iteration, i.e., $F^e(r) = r^q$. For an ideal $I = (x_1, \dots, x_n) \subseteq R$, we let $I^{[q]} = (x_1^q, \dots, x_n^q)$. Note that $F^e(I)R = I^{[q]}$, where $q = p^e$, as always.

We shall denote by R° the complement of the union of the minimal primes of R .

Date: October 16, 1997.

Definition 2.1. A ring R is said to be F -pure if the Frobenius homomorphism $F : M \rightarrow M \otimes_R F(R)$ is injective for all R -modules M .

For an element x of R and an ideal I , we say that $x \in I^F$, the *Frobenius closure* of I , if there exists $q = p^e$ such that $x^q \in I^{[q]}$. A normal domain R is F -pure if and only if for all ideals I of R , we have $I^F = I$.

We say that $x \in I^*$, the *tight closure* of I , if there exists $c \in R^\circ$ such that $cx^q \in I^{[q]}$ for all $q = p^e \gg 0$.

It is easily verified that $I \subseteq I^F \subseteq I^*$. Furthermore, I^* is always contained in the integral closure of I and is frequently much smaller.

3. PRELIMINARY COMPUTATIONS

We record some determinant computations we shall find useful. Note that for integers n and m where $m \geq 1$, we shall use the notation:

$$\binom{n}{m} = \frac{(n)(n-1)\cdots(n-m+1)}{(m)(m-1)\cdots(1)}.$$

Lemma 3.1.

$$\det \begin{vmatrix} \binom{n}{a+k} & \binom{n}{a+k+1} & \cdots & \binom{n}{a+2k} \\ \binom{n}{a+k-1} & \binom{n}{a+k} & \cdots & \binom{n}{a+2k-1} \\ \cdots & \cdots & \cdots & \cdots \\ \binom{n}{a} & \binom{n}{a+1} & \cdots & \binom{n}{a+k} \end{vmatrix} = \frac{\binom{n}{a+k} \binom{n+1}{a+k} \cdots \binom{n+k}{a+k}}{\binom{a+k}{a+k} \binom{a+k+1}{a+k} \cdots \binom{a+2k}{a+k}}.$$

Proof. This is evaluated in [Mu, page 682] as well as [Ro]. □

Lemma 3.2. Let $F(n, a, k)$ denote the determinant of the matrix

$$M(n, a, k) = \begin{pmatrix} \binom{n}{a} & \binom{n}{a+1} & \cdots & \binom{n}{a+k} \\ \binom{n+2}{a+1} & \binom{n+2}{a+2} & \cdots & \binom{n+2}{a+k+1} \\ \cdots & \cdots & \cdots & \cdots \\ \binom{n+2k}{a+k} & \binom{n+2k}{a+k+1} & \cdots & \binom{n+2k}{a+2k} \end{pmatrix}.$$

Then for $k \geq 1$ we have

$$\frac{F(n, a, k)}{F(n+2, a+2, k-1)} = \binom{n}{a} \prod_{s=1}^k \prod_{r=1}^k \frac{s(s+2a-n)}{(a+r)(n-a+r)}.$$

Hence

$$F(n, a, k) = \frac{\binom{n}{a} \binom{n+2}{a+2} \cdots \binom{n+2k}{a+2k}}{\binom{a+k}{k} \binom{a+k+1}{k-1} \cdots \binom{a+2k-1}{1}} \cdot \frac{\binom{2a-n+k}{k} \binom{2a-n+k+1}{k-1} \cdots \binom{2a-n+2k-1}{1}}{\binom{n-a+k}{k} \binom{n-a+k-1}{k-1} \cdots \binom{n-a+1}{1}}.$$

Proof. We shall perform row operations on $M(n, a, k)$ in order to get zero entries in the first column from the second row onwards, starting with the last row and moving up. More precisely, from the $(r+1)$ th row, subtract the r th row multiplied by

$\binom{n+2r}{a+r}/\binom{n+2r-2}{a+r-1}$ starting with $r = k$, and continuing until $r = 2$. The $(r+1, s+1)$ th entry of the new matrix, for $r \geq 1$, is

$$\binom{n+2r}{a+r+s} - \frac{\binom{n+2r}{a+r}}{\binom{n+2r-2}{a+r-1}} \binom{n+2r-2}{a+r-1+s} = \frac{s(s+2a-n)}{(a+r)(n-a+r)} \binom{n+2r}{a+r+s}.$$

We have only one nonzero entry in the first column, namely $\binom{n}{a}$ and so we examine the matrix obtained by deleting the first row and column. Factoring out $s(s+2a-n)$ from each column for $s = 1, \dots, k$ and $1/(a+r)(n-a+r)$ from each row for $r = 1, \dots, k$, we see that

$$\det M(n, a, k) = \binom{n}{a} \prod_{s=1}^k \prod_{r=1}^k \frac{s(s+2a-n)}{(a+r)(n-a+r)} \det M(n+2, a+2, k-1).$$

The required result immediately follows. $\square$

Lemma 3.3. *Consider the polynomial ring $T = K[A_1, \dots, A_m]$ where $I_{r,i}$ denotes the ideal $I_{r,i} = (A_1^i, \dots, A_r^i)T$ for $r \leq m$. Then*

$$(A_1 \cdots A_{r-1})^\alpha (A_1 + \cdots + A_{r-1})^\beta \in I_{r-1, \alpha+\gamma} + (A_1 + \cdots + A_{r-1})^{\alpha+\gamma} T$$

for positive integers α, β , and γ implies

$$(A_1 \cdots A_r)^\alpha (A_1 + \cdots + A_r)^{\beta+\gamma-1} \in I_{r, \alpha+\gamma} + (A_1 + \cdots + A_r)^{\alpha+\gamma} T.$$

Proof. Consider the binomial expansion of $(A_1 \cdots A_r)^\alpha (A_1 + \cdots + A_r)^{\beta+\gamma-1}$ into terms of the form $(A_1 \cdots A_{r-1})^\alpha (A_1 + \cdots + A_{r-1})^{\beta+\gamma-1-j} A_r^{\alpha+j}$. Such an element is clearly in $I_{r, \alpha+\gamma}$ whenever $j \geq \gamma$, and so assume $\gamma > j$. Now

$$\begin{aligned} (A_1 \cdots A_{r-1})^\alpha A_r^{\alpha+j} (A_1 + \cdots + A_{r-1})^{\beta+\gamma-1-j} \\ \in I_{r, \alpha+\gamma} + A_r^{\alpha+j} (A_1 + \cdots + A_{r-1})^{\alpha+2\gamma-1-j} T \\ \subseteq I_{r, \alpha+\gamma} + (A_1 + \cdots + A_{r-1}, A_r)^{2\alpha+2\gamma-1} T \\ \subseteq I_{r, \alpha+\gamma} + (A_1 + \cdots + A_r)^{\alpha+\gamma} T. \end{aligned}$$

$\square$

4. TIGHT CLOSURE

We now prove the main theorem.

Theorem 4.1. *Let $R = K[X_1, \dots, X_n]/(X_1^n + \cdots + X_n^n)$ where $n \geq 3$ and K is a field of prime characteristic p where $p \nmid n$. Then*

$$(x_1 \cdots x_n)^{n-2} \in (x_1^{n-1}, \dots, x_n^{n-1})^*.$$

Note that there are infinitely many $e \in \mathbb{N}$ such that $p^e = q \equiv 1 \pmod{n}$. By [HH, Lemma 8.16], it suffices to work with powers of p of this form, and show that for all such q we have

$$(x_1 \cdots x_n)^{(n-2)q+1} \in (x_1^{(n-1)q}, \dots, x_n^{(n-1)q}).$$

Letting $q = nk + 1$, it suffices to show

$$(x_1 \cdots x_n)^{(n-2)nk} \in (x_1^{(n-1)nk}, \dots, x_n^{(n-1)nk}).$$

Let $A_1 = x_1^n, \dots, A_n = x_n^n$ and note that $A_1 + \dots + A_n = 0$. In this notation, we aim to show

$$(A_1 \cdots A_n)^{(n-2)k} \in (A_1^{(n-1)k}, \dots, A_n^{(n-1)k}).$$

Our task is then effectively reduced to working in the polynomial ring

$$K[A_1, \dots, A_{n-1}] \cong K[A_1, \dots, A_n]/(A_1 + \dots + A_n)$$

where we need to show

$$(A_1 \cdots A_{n-1}(A_1 + \cdots + A_{n-1}))^{(n-2)k} \in I_{n-1, (n-1)k} + (A_1 + \cdots + A_{n-1})^{(n-1)k}.$$

By repeated use of Lemma 3.3, it suffices to show

$$(A_1 A_2)^{(n-2)k} (A_1 + A_2)^k \in (A_1^{(n-1)k}, A_2^{(n-1)k}, (A_1 + A_2)^{(n-1)k}).$$

We have now reduced our problem to a statement about a polynomial ring in two variables. The required result follows from the next lemma.

Lemma 4.2. *Let $K[A, B]$ be a polynomial ring over a field K of characteristic $p > 0$ and e be a positive integer such that $q = p^e \equiv 1 \pmod{n}$. If $q = nk + 1$, we have*

$$(A, B)^{(2n-3)k} \subseteq I = (A^{(n-1)k}, B^{(n-1)k}, (A+B)^{(n-1)k}).$$

In particular, $(AB)^{(n-2)k}(A+B)^k \in I$.

Proof. Note that I contains the following elements: $(A+B)^{(n-1)k} A^k B^{(n-3)k}$, $(A+B)^{(n-1)k} A^{k-1} B^{(n-3)k+1}, \dots, (A+B)^{(n-1)k} B^{(n-2)k}$. We take the binomial expansions of these elements and consider them modulo the ideal $(A^{(n-1)k}, B^{(n-1)k})$. This shows that the following elements are in I :

$$\begin{array}{ccccccc} \binom{(n-1)k}{k} A^{(n-1)k} B^{(n-2)k} & + & \cdots & + & \binom{(n-1)k}{2k} A^{(n-2)k} B^{(n-1)k}, \\ \binom{(n-1)k}{k-1} A^{(n-1)k} B^{(n-2)k} & + & \cdots & + & \binom{(n-1)k}{2k-1} A^{(n-2)k} B^{(n-1)k}, \\ \cdots & & & & \\ \hline \binom{(n-1)k}{0} A^{(n-1)k} B^{(n-2)k} & + & \cdots & + & \binom{(n-1)k}{k} A^{(n-2)k} B^{(n-1)k}. \end{array}$$

The coefficients of $A^{(n-1)k}B^{(n-2)k}, A^{(n-1)k-1}B^{(n-2)k+1}, \dots, A^{(n-2)k}B^{(n-1)k}$ form the matrix:

$$\begin{pmatrix} \binom{(n-1)k}{k} & \binom{(n-1)k}{k+1} & \cdots & \binom{(n-1)k}{2k} \\ \binom{(n-1)k}{k-1} & \binom{(n-1)k}{k} & \cdots & \binom{(n-1)k}{2k-1} \\ \vdots & \vdots & \ddots & \vdots \\ \binom{(n-1)k}{0} & \binom{(n-1)k}{1} & \cdots & \binom{(n-1)k}{k} \end{pmatrix}.$$

To show that all monomials of degree $(2n-3)k$ in A and B are in I , it suffices to show that this matrix is invertible. Since $q = nk + 1$ we have $\binom{(n-1)k+r}{k} = (-1)^k \binom{2k-r}{k}$ for $0 \leq r \leq k$, and so by Lemma 3.1, the determinant of this matrix is

$$\frac{\binom{(n-1)k}{k} \binom{(n-1)k+1}{k} \cdots \binom{nk}{k}}{\binom{k}{k} \binom{k+1}{k} \cdots \binom{2k}{k}} = (-1)^{k(k+1)} \frac{\binom{2k}{k} \binom{2k-1}{k} \cdots \binom{k}{k}}{\binom{k}{k} \binom{k+1}{k} \cdots \binom{2k}{k}} = 1.$$



With this we complete the proof that $(x_1 \cdots x_n)^{n-2} \in (x_1^{n-1}, \dots, x_n^{n-1})^*$.

5. FROBENIUS CLOSURE

Let $R = K[X_1, \dots, X_n]/(X_1^n + \dots + X_n^n)$ as before, where the characteristic of K is $p \nmid n$.

Lemma 5.1. *Let $R = K[X_1, \dots, X_n]/(X_1^n + \dots + X_n^n)$ where K is a field of characteristic p . Then R is F -pure if and only if $p \equiv 1 \pmod{n}$.*

Proof. This is Proposition 5.21 (c) of [HR]. $\square$

The main result of this section is the following theorem.

Theorem 5.2. *Let $R = K[X_1, \dots, X_n]/(X_1^n + \dots + X_n^n)$ where K is a field of characteristic p . Then*

$$(x_1 \cdots x_n)^{n-2} \in (x_1^{n-1}, \dots, x_n^{n-1})^F.$$

if and only if $p \not\equiv 1 \pmod{n}$.

One implication follows from Lemma 5.1, and so we need to consider the case $p \not\equiv 1 \pmod{n}$.

The case $n = 3$ seems to be the most difficult, and we handle that first. Let $R = K[X, Y, Z]/(X^3 + Y^3 + Z^3)$ where $p \equiv 2 \pmod{3}$. We need to show that $xyz \in (x^2, y^2, z^2)^F$.

Let $A = y^3, B = z^3$ and so $A + B = -x^3$. We first show that when $p = 2$, we have $xyz \in (x^2, y^2, z^2)^F$ by establishing that $(xyz)^8 \in (x^2, y^2, z^2)^{[8]}$. Note that it suffices to show that $(xyz)^6 \in (x^{15}, y^{15}, z^{15})$, or in other words that $(AB(A+B))^2 \in (A^5, B^5, (A+B)^5)$, but this is easily seen to be true.

We may now assume $p = 6m + 5$ where $m \geq 0$. We shall show that in this case $(xyz)^p \in (x^2, y^2, z^2)^{[p]}$, i.e., that

$$(xyz)^{6m+5} \in (x^{12m+10}, y^{12m+10}, z^{12m+10}).$$

Note that to establish this, it suffices to show

$$(xyz)^{6m+3} \in (x^{12m+9}, y^{12m+9}, z^{12m+9}),$$

i.e., that $(AB(A+B))^{2m+1} \in (A^{4m+3}, B^{4m+3}, (A+B)^{4m+3})$.

Lemma 5.3. *Let $K[A, B]$ be a polynomial ring over a field K of characteristic $p = 6m + 5$ where $m \geq 0$. Then we have*

$$(AB(A+B))^{2m+1} \in I = (A^{4m+3}, B^{4m+3}, (A+B)^{4m+3}).$$

Proof. To show that $(AB(A+B))^{2m+1} \in I$, we shall show that the following terms grouped together symmetrically from its binomial expansion,

$$\begin{aligned} f_1 &= (AB)^{3m+1}(A+B), \quad f_3 = (AB)^{3m}(A^3 + B^3), \quad \dots, \\ f_{2m+1} &= (AB)^{2m+1}(A^{2m+1} + B^{2m+1}), \end{aligned}$$

are all in the ideal I . Note that I contains the elements $(AB)^m(A+B)^{4m+3}$, $(AB)^{m-1}(A+B)^{4m+5}$, $\dots$, $(AB)(A+B)^{6m+1}$, $(A+B)^{6m+3}$. We consider the binomial expansions of these elements modulo (A^{4m+3}, B^{4m+3}) , and get the following

$$\begin{array}{ccccccc} \binom{4m+3}{2m+2} f_1 & + & \binom{4m+3}{4m+5} f_3 & + & \dots & + & \binom{4m+3}{3m+2} f_{2m+1}, \\ \binom{4m+5}{2m+3} f_1 & + & \binom{4m+5}{2m+4} f_3 & + & \dots & + & \binom{4m+5}{3m+3} f_{2m+1}, \\ & & & & \dots & & \\ & & & & \dots & & \\ & & & & \dots & & \\ \binom{6m+3}{3m+2} f_1 & + & \binom{6m+3}{3m+3} f_3 & + & \dots & + & \binom{6m+3}{4m+2} f_{2m+1}. \end{array}$$
$$\begin{pmatrix} \binom{4m+3}{2m+2} & \binom{4m+3}{2m+3} & \cdots & \binom{4m+3}{3m+2} \\ \binom{4m+5}{2m+3} & \binom{4m+5}{2m+4} & \cdots & \binom{4m+5}{3m+3} \\ \vdots & \vdots & \ddots & \vdots \\ \binom{6m+3}{3m+2} & \binom{6m+3}{3m+3} & \cdots & \binom{6m+3}{4m+2} \end{pmatrix}.$$
$$(x_1 \cdots x_n)^{(n-2)p} = -(x_1 \cdots x_{n-1})^{(n-2)p} x_n^{(n-2)p-n} (x_1^n + \cdots + x_{n-1}^n) \\ \in (x_1^{(n-1)p}, \dots, x_{n-1}^{(n-1)p}).$$
$$(x_1 \cdots x_n)^{(n-2)p} \in (x_1^{(n-1)p}, \dots, x_n^{(n-1)p}).$$
$$(x_1 \cdots x_n)^{(n-2)nk} \in (x_1^{(n-1)nk+n}, \dots, x_n^{(n-1)nk+n}).$$
$$(A_1 \cdots A_n)^{(n-2)k} \in (A_1^{(n-1)k+1}, \dots, A_n^{(n-1)k+1}).$$
$$(A_1 A_2)^{(n-2)k} (A_1 + A_2)^k \in I = (A_1^{(n-1)k+1}, A_2^{(n-1)k+1}, (A_1 + A_2)^{(n-1)k+1}).$$
$$(A, B)^{(2n-3)k} \subseteq I = (A^{(n-1)k+1}, B^{(n-1)k+1}, (A+B)^{(n-1)k+1}).$$
$$(A+B)^{(n-1)k+1}A^{k-1}B^{(n-3)k}, \dots, (A+B)^{(n-1)k+1}B^{(n-2)k-1}.$$

$$\begin{array}{ccccccc} \binom{(n-1)k+1}{k+1} A^{(n-1)k} B^{(n-2)k} & + & \cdots & + & \binom{(n-1)k+1}{2k+1} A^{(n-2)k} B^{(n-1)k}, \\ \binom{(n-1)k+1}{k} A^{(n-1)k} B^{(n-2)k} & + & \cdots & + & \binom{(n-1)k+1}{2k} A^{(n-2)k} B^{(n-1)k}, \\ \hline \binom{(n-1)k+1}{1} A^{(n-1)k} B^{(n-2)k} & + & \cdots & + & \binom{(n-1)k+1}{k+1} A^{(n-2)k} B^{(n-1)k}. \end{array}$$
$$\begin{pmatrix} \binom{(n-1)k+1}{k+1} & \binom{(n-1)k+1}{k+2} & \cdots & \binom{(n-1)k+1}{2k+1} \\ \binom{(n-1)k+1}{k} & \binom{(n-1)k+1}{k+1} & \cdots & \binom{(n-1)k+1}{2k} \\ \vdots & \vdots & \ddots & \vdots \\ \binom{(n-1)k+1}{1} & \binom{(n-1)k+1}{2} & \cdots & \binom{(n-1)k+1}{k+1} \end{pmatrix}.$$
$$\frac{\binom{(n-1)k+1}{k+1} \binom{(n-1)k+2}{k+1} \cdots \binom{nk+1}{k+1}}{\binom{k+1}{k+1} \binom{k+2}{k+1} \cdots \binom{2k+1}{k+1}}$$
$$R = K[X, Y, Z]/(X^3 + Y^3 + Z^3)$$
$$(zu)(zv)(xv - yu) = (zv)^2(xu) - (zu)^2(yv).$$
$$(zu)(zv) \in \quad (xu, yv) :_T (xv - yu) \quad \subseteq (xu, yv)^*.$$

There is a retraction $R \otimes_K S \rightarrow R$ under which $U \mapsto 1$ and $V \mapsto 1$. This gives us a retraction from $T \rightarrow R$ which, when applied to $(zu)(zv) \in (xu, yv)^*$, shows $z^2 \in (x, y)^*$ in R .

REFERENCES

- [HH] M. Hochster and C. Huneke, Tight closure, invariant theory, and the Briançon-Skoda theorem, *J. Amer. Math. Soc.* **3** (1990), 31–116.
- [Ho] M. Hochster, Tight closure in equal characteristic, big Cohen–Macaulay algebras, and solid closure, *Contemp. Math.* **159** (1994), 173–196.
- [HR] M. Hochster and J. Roberts, The purity of the Frobenius and local cohomology, *Adv. in Math.* **21** (1976), 117–172.
- [Hu] C. Huneke, Tight closure, parameter ideals and geometry, in “Commutative Algebra,” Six lectures on commutative algebra (Bellaterra, 1996), 187–239, Progr. Math., 166, Birkhäuser, Basel, 1998.
- [Mc] M. McDermott, “Tight closure, plus closure and Frobenius closure in cubical cones,” Thesis, Univ. of Michigan, 1996.
- [Mu] T. Muir, “A treatise on the theory of determinants,” Revised and enlarged by W. H. Metzler, Longmans, Green and co., New York, London, 1933.
- [Ro] P. Roberts, A computation of local cohomology, *Contemp. Math.* **159** (1994), 351–356.
- [Sm] K. E. Smith, Tight Closure of parameter ideals, *Invent. Math.* **115** (1994), 41–60.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF MICHIGAN, EAST HALL, 525 EAST UNIVERSITY AVENUE, ANN ARBOR, MI 48109-1109

E-mail address: `binku@math.lsa.umich.edu`